Mon. Not. R. Astron. Soc. **000**, 1–7 (2015) Printed 29 April 2016 (MN LATEX style file v2.2)

Stark broadening of B IV spectral lines

Milan S. Dimitrijević[1,2,3,4⋆], Magdalena Christova[5], Zoran Simić[1], Andjelka Kovačević[6] and Sylvie Sahal-Bréchot[3]

[1] *Astronomical Observatory, Volgina 7, 11060 Belgrade, Serbia.*
[2] *IHIS -Techno Experts, Bežanijski put 23, 11080 Belgrade-Zemun, Serbia.*
[3] *LERMA, Observatoire de Paris, PSL Research University, CNRS, Sorbonne Universities, UPMC Univ. Paris 06, 5 Place Jules Janssen, 92195 Meudon Cedex, France.*
[4] *Institute Isaac Newton of Chile, Yugoslavia Branch, 11060 Belgrade, Serbia.*
[5] *Department of Applied Physics, Technical University-Sofia, 1000 Sofia, Bulgaria.*
[6] *Department for Astronomy, Faculty for Mathematics, Studentski Trg 16, 11000 Belgrade, Serbia.*

arXiv:1604.08537v1 [astro-ph.SR] 28 Apr 2016

29 April 2016

ABSTRACT

Stark broadening parameters for 157 multiplets of helium like boron (B IV) have been calculated using the impact semiclassical perturbation formalism. Obtained results have been used to investigate the regularities within spectral series. An example of the influence of Stark broadening on B IV lines in DO white dwarfs is given.

Key words: atomic data – atomic processes – line: formation

1 INTRODUCTION

The study presents Stark broadening parameters (widths and shifts) of B IV spectral lines calculated using the impact semi-classical perturbation formalism (Sahal-Bréchot 1969a,b). B IV is a helium-like ion and such ions are rather commonly used for the diagnostic (Kolk, König & Kunze 1986) of laboratory (TFR Group, Doyle & Schwob, 1982; Boiko et al. 1983; Källne, Källne & Pradhan 1983) and astrophysical plasmas (McKenzie & Landecker 1982).

In astrophysics, Stark broadening data for various atomic and ionic lines (since we do not know *a priori* the chemical composition of a star) are of particular interest especially for white dwarfs, where this line broadening mechanism is usually the principal one (Popović et al. 1999b; Tankosić, Popović & Dimitrijević 2003; Milovanović et al. 2004; Simić et al. 2006; Hamdi et al. 2008; Dimitrijević et al. 2011; Dufour et al. 2011; Larbi-Terzi et al. 2012; Simić, Dimitrijević & Sahal-Bréchot 2013; Simić, Dimitrijević & Popović 2014). This broadening mechanism may be of interest and for the main sequence stars, especialy for A type and late B type (Lanz et al. 1988; Popović et al 1999a; Popović et al. 1999b; Popović et al 2001a,b; Dimitrijević et al. 2003a,b; Tankosić, Popović & Dimitrijević 2003; Dimitrijević et al. 2004; Milovanović et al. 2004; Dimitrijević et al. 2005; Simić et al. 2005a,b; Simić, Dimitrijević & Kovačević 2009; Simić, Dimitrijević & Sahal-Bréchot 2013; Simić, Dimitrijević & Popović 2014).

We note as well the increasing astrophysical importance of Stark broadening data for various atoms and ions of trace elements, without astrophysical meaning before the development of satellite born telescopes which now are providing high-resolution spectra of earlier inaccessible quality. Well-resolved line profiles for many white dwarfs, where Stark broadening is important, have been and will be provided for example by the Space Telescope Imaging Spectrograph (STIS), Cosmic Origins Spectrograph (COS) and Goddard High Resolution Spectrograph (GHRS), Far Ultraviolet Spectroscopy Explorer (FUSE), the International Ultraviolet Explorer and others.

Data on boron lines, including Stark broadening, are of interest in astrophysics but also for example for laboratory (Blagojević et al. 1999), fusion (Iglesias et al. 1998) and laser produced (Nicolosi et al. 1978) plasmas investigations as well as for laser research and development (Wang et al. 1992). In astrophysics, the light elements lithium, beryllium, and boron are of great interest for two sets of reasons, which might be categorized as cosmological and related to stellar structure (Duncan et al. 1998). A general trend in nature is that the abundance of the elements versus the mass number draws a globally decreasing curve (Vangioni-Flam & Cassé 1999; Vangioni-Flam, Cassé & Audouze 2000). However, the rare and fragile light nuclei, lithium, beryllium and boron are not generated in the normal course of stellar nucleosynthesis (except ^{7}Li, in the galactic disk) and are, in fact, destroyed in stellar interiors. The standard Big Bang nucleosynthesis (BBN) theory is not effective to explain the generation of ^{6}Li, ^{9}Be, ^{10}B, ^{11}B (Delbourgo-Salvador & Vangioni-Flam 1993; Schramm 1993; Thomas et al. 1993), what is reflected in the low abundance of these simple species. Consequently

⋆ E-mail: mdimitrijevic@aob.bg.ac.rs

the origin and evolution of boron, are of particular interest and the corresponding Stark broadening data are needed (Tankosić, Popović & Dimitrijević 2003).

The importance of light element abundance for the giant-branch evolution is underlined in Duncan et al. (1998). The stellar structure interest steems from the fact that Li, Be and B undergo nuclear reactions at relatively low temperatures, approximately 2.5, 3.5, and 5 $\times 10^6$ K at densities similar to those in the Sun. Since these temperatures are reached not far below the convection zone and well outside the core in solar-type stars, circulation and destruction of the light elements can result in observable abundance changes. Observations of these changes can provide an invaluable probe of stellar structure and mixing. Both Li and Be abundances are greatly reduced in the giants from their initial main-sequence values. Duncan et al. (1998) report the B abundance of two giants and one dwarf in the Hyades, the latter included to evaluate explicitly the boron abundance prior to giant-branch evolution. They demonstrate empirically that boron contributes to the absorption spectra of cool stars. HST measurements of boron abundances of these objects have permitted a test of one of the basic predictions of stellar evolution theory: the growth of the convection zone as a star evolves up the giant branch.

Cunha, Smith & Lambert (1999) report on the base of Hubble Space Telescope Goddard High Resolution Spectrograph spectra, the boron abundance derived for the young Orion solar-type member BD - 05°1317. They note that, the real interstellar boron abundance and its comparison with the stellar values remains uncertain. Determinations of the Orion associations boron abundance provide unexpected results. The boron abundance derived from spectra of B-type stars is consistent with the expectation that the boron abundance of the Orion association should be similar to that of the solar system, but is considerably higher than the interstellar boron abundance for several lines of sight, including some toward Orion. A low boron abundance raises the question as to how the boron abundance of interstellar gas and young stars has decreased by a factor of 4 or 5 since the solar system was formed Cunha, Smith & Lambert (1999).

The light trace elements lithium, beryllium, and boron are at the center of astrophysical puzzles involving topics as diverse as the primordial fireball, interstellar (IS) or even intergalactic space, and stellar surfaces and interiors (Venn et al. 2002). This role arises because boron nuclei are destroyed by warm protons, and thus even quite shallow mixing of the atmosphere with the interior reduces the surface abundance by bringing boron-depleted material to the surface. In Venn et al. (2002), a study on boron abundance of B-type stars has been presented. Boron alone is observable in hot stars where Stark broadening is not negligible or important. A principal goal of most of these studies of hot stars was to establish the present-day boron abundance in order to improve our understanding of the Galactic chemical evolution of boron. Boron in hot stars, like lithium in cool stars, is shown to be a tracer of some of the various processes afecting a stars surface composition that are not included in the standard models of stellar evolution. If the initial boron abundances of local hot stars are similar from star to star and association to association, then the large spread in boron abundances, a factor of at least 30 across their sample, shows that boron abundances are a clue to unraveling the nonstandard processes that afect young hot stars. We note that in hot stars Stark broadening is often needed for the determination of abundances and in Popović et al. (1999b) are analysed errors in abundances if it is not taken into account, especially for A-type stars.

Spectral lines of the boron ions have been observed in stellar spectra. B I lines have been observed in F and G stars (Duncan et al. 1997), B II in hotter stars (Cunha et al. 1997) and B III in early B stars (Cunha et al. 1997; Proffitt et al. 1999). For example, Proffitt & Quigley (2001) found B III lines in 44 early B stars.

Recently, we have calculated Stark broadening parameters for 36 B IV multiplets (Dimitrijević et al. 2014). To complete as much as possible the corresponding Stark broadening data needed in astrophysics, laboratory-, technological-, fusion-, and laser produced-plasma physics, our aim is to present in this work new theoretical determinations of Stark broadening parameters (full widths at half intensity and shifts) within the impact semiclassical perturbation approach for additional 121 helium like boron (B IV) multiplets, so that that semiclassical perturbation Stark widths for 157 B IV multiplets will be available. Moreover, in Dimitrijević et al. (2014) data are provided only for an electron density of $N = 10^{17}$ cm^{-3}, while here, Stark broadening parameters are calculated for wide range of perturber densities. Obtained results will be used for the consideration of regulariries within spectral series, as well as for one example of the influence of Stark broadening in comparison with Doppler broadening on B IV lines in DO white dwarfs.

2 THE IMPACT SEMICLASSICAL PERTURBATION METHOD

Pressure broadening of spectral lines arises when an atom, ion, or molecule which emits or absorbs light in a gas or plasma, is perturbed by its interactions with the other particles of the medium. Interpretation of this phenomenon is currently used for modelling of the medium and for spectroscopic diagnostics, since the broadening of the lines depends on the temperature and density of the medium. The physical conditions in the Universe are very various, and collisional broadening with charged particles (Stark broadening) appears to be important in many domains. For example, at temperatures around 10^4 K and densities 10^{13} - 10^{15} cm^{-3}, Stark broadening is of interest for modelling and analysing spectra of A and B type stars (see e.g. Lanz et al. (1988); Popović et al (1999a); Popović et al. (1999b); Popović et al (2001a,b); Tankosić, Popović & Dimitrijević (2003); Simić et al. (2005a,b)). In white dwarfs, especially Stark broadening is the dominant collisional line broadening mechanism in all important layers of the atmosphere (Popović et al. 1999b; Tankosić, Popović & Dimitrijević 2003; Milovanović et al. 2004; Simić et al. 2006; Dimitrijević et al. 2011; Dufour et al. 2011; Larbi-Terzi et al. 2012; Simić, Dimitrijević & Sahal-Bréchot 2013; Simić, Dimitrijević & Popović 2014). The theory of Stark broadening is well applied, especially for accurate spectroscopic diagnostics and modelling. This requires the knowledge of numerous profiles, especially for trace elements, as boron in this case, which are used as useful probes for modern spectroscopic diagnostics. Interpretation

of the spectra of white dwarfs, which are very faint, allows understanding the evolution of these very old stars, which are close to death.

The results for Stark broadening parameters of helium like boron (B IV) multiplets have been calculated using the semi-classical perturbation formalism (Sahal-Bréchot 1969a,b).We note that different later inovations and optimizations of this theoretical method have been described in details in Sahal-Bréchot (1974, 1991); Fleurier, Sahal-Bréchot & Chapelle (1977); Dimitrijević, Sahal-Bréchot & Bommier (1991); Dimitrijević & Sahal-Bréchot (1996); Sahal-Bréchot et al. (2014). Within this theory the FWHM - Full Width at Half intensity Maximum (W) and the shift (d) of an isolated line, originating from the transition between the initial level i and the final level f is expressed for an ionized atom as:

$$W = N\int vf(v)dv\left(\sum_{i'\neq i}\sigma_{ii'}(v)+\sum_{f'\neq f}\sigma_{ff'}(v)+\sigma_{el}\right)$$

$$d = N\int vf(v)dv\int_{R_3}^{R_D}2\pi\rho d\rho\sin(2\varphi_p). \quad (1)$$

where i' and f' are perturbing levels, N and v are the electron density and the velocity of perturbers respectively, $f(v)$ is the Maxwellian distribution of electron velocities, and ρ the impact parameter of the free electron colliding with the emitter.

The inelastic cross sections $\sigma_{ii'}(v)$ (respectively $\sigma_{ff'}(v)$), can be expressed by an integration of the transition probability $P_{jj'}(\rho,v), j=i,f; j'=i',f'$, over the impact parameter ρ as:

$$\sum_{i'\neq i}\sigma_{ii'}(v)=\frac{1}{2}\pi R_1^2+\int_{R_1}^{R_D}2\pi\rho d\rho\sum_{i'\neq i}P_{ii'}(\rho,v). \quad (2)$$

and the elastic contribution to the width is given by:

$$\sigma_{el}=2\pi R_2^2+\int_{R_2}^{R_D}2\pi\rho d\rho\sin^2\delta+\sigma_r,$$

$$\delta=(\varphi_p^2+\varphi_q^2)^{\frac{1}{2}}. \quad (3)$$

Here σ_{el} is the elastic cross section, while φ_p (r^{-4}) and φ_q (r^{-3}), are phase shifts due to the polarization and quadrupolar potential respectively, and are defined in Section 3 of Chapter 2 in Sahal-Bréchot (1969a). The cut-offs R_1, R_2, R_3, and the Debye radius R_D, as well as the symmetrization procedure are described in Section 1 of Chapter 3 in Sahal-Bréchot (1969b). The contribution of Feshbach resonances, σ_r is explained in Fleurier, Sahal-Bréchot & Chapelle (1977).

Within the semiclassical perturbation theory is assumed that electrons are moving along hyperbolic paths due to attractive Coulomb force, while in the case of ionic perturbers this force is repulsive, so that trajectories are different. For ion-impact broadening the formulae are analogous to Eqs. (1) - (3) for electron-impact broadening, but in the case of ion broadening there is no term for the contribution of Feshbach resonances.

If the considered lines are isolated, the line profile $F(\omega)$ has Lorentzian form and can be expressed as:

$$F(\omega)=\frac{W/(2\pi)}{(\omega-\omega_{if}-d)^2+(W/2)^2} \quad (4)$$

where

$$\omega_{if}=\frac{E_i-E_f}{\hbar}$$

and E_i, E_f are the energies of initial and final state, respectively. Consequently, if we know Stark width W and shift d it is easy to determine the profile of a considered spectral line.

3 STARK BROADENING PARAMETER CALCULATIONS

We have calculated here, within the frame of semiclassical perturbation method (Sahal-Bréchot 1969a,b), electron-, proton-, and helium ion-impact broadening parameters, full width at half maximum of intensity (FWHM - W) and shift (d) for 157 transitions of B IV, for temperatures of 50 000K, 100 000 K, 200 000 K and 500 000K and perturber densities from 10^{14} to 10^{22} cm^{-3}. Energy levels needed for these calculations, have been taken from Kramida et al. (2008), while for the needed oscillator strengths Bates & Damgaard (1949) method has been used, together with the tables of Oertel & Shomo (1968) and the article of Van Regemorter, Hoang Binh & Prud'homme (1979) for higher levels.

We note here, that experimental lifetimes for levels corresponding to 385.0 Å and 1170Å can be found in Kernahan et al. (1975). However, for semiclassical perturbation calculations of Stark broadening parameters we need a consistent set of oscillator strengths and if we use a mixture of oscillator strengths obtained with various methods this may result in worse agreement with experimental data for Stark broadening, as shown in Dimitrijević & Sahal-Bréchot (1994), since the corresponding summation rule for oscillator strengts for transitions starting or ending on a particular energy level, may be violated.

The complete results for all 157 transitions, are provided in electronic, computer readable form, as additional data in the on line journal editionas as Tables S1-S7. In Table S1 are results for 10^{14} cm^{-3} and in Tables S2-S7 for 10^{17} to 10^{22} cm^{-3}. As an example, here are given in Table 1, the results for two transitions as a sample in order to demonstrate the form of additional data.

It should be noticed that wavelengths in Table 1 as well as in Tables S1-S7 differs from experimental ones since they are calculated. Since the Stark broadening parameter calculation depends only on relative positions of energy levels they do not depend on absolute positions of energy levels when expressed in angular frequency units. The Stark width in Å may be expressed in angular frequency units using the formula:

$$W(\mathring{A})=\frac{\lambda^2}{2\pi c}W(s^{-1}) \quad (5)$$

where c is the speed of light. If a correction of the width or

shift for the difference between calculated and experimental wavelength is necessary this can be made for the width, and similarly for the shift as:

$$W_{cor} = \left(\frac{\lambda_{\rm exp}}{\lambda}\right)^2 W. \quad (6)$$

Here, with W_{cor} is denoted the corrected width, λ_{exp} is the experimental, λ the calculated wavelength and W the width from Table 1, or Tables S1-S7.

A parameter C (Dimitrijević & Sahal-Bréchot 1984), provided in Table 1 and Tables 1S-7S, enables to determine the maximal perturber density when the line may be considered as isolated, by division of C with the corresponding full width at half maximum. The validity of impact approximation is checked for all results estimating the value of NV, where V is the collision volume and N the perturber density. If $NV<$ 0.1, we assume that the impact approximation is valid (Sahal-Bréchot 1969a,b). If $NV>$ 0.5 the corresponding results for width and shift are not shown in tables since the impact approximation breaks down, while for $0.1<NV\leqslant 0.5$ befor the corresponding Stark broadening parameters is an asterisk to notice that this result is on the limit of validity of impact approximation. In the cases when, for the given (T and N), the impact approximation is not valid, for the estimation of ion broadening is convenient quasistatic approach (Griem 1974; Sahal-Bréchot 1991). If neither impact nor quasistatic approximation are valid, one can use a unified-type theory, as for example the approach of Barnard, Cooper & Smith (1974) providing a simple analytical formula, convenient to use for such a case.

The dependence of the broadening parameters of spectral lines due to impacts with charged particles versus principal quantum number within a spectral series is a useful information. If we know the trend of Stark broadening parameters within a spectral series, it is possible to interpolate or extrapolate the eventually missing values within the considered series. The regular behavior of electron Stark broadening width versus principal quantum number within three spectral series, 1s^2 ^{1}S - 1snp ^{1}P, 1s2s ^{1}S - 1snp ^{1}P and 1s3p ^{1}P - 1snd ^{1}D is presented in Fig. 1. The widths increase with principal quantum number which reflects the decrease of the distance to the nearest perturbing levels with the increase of the principal quantum number.

The calculated data for 1s^2 ^{1}S - 1snp ^{1}P and 1s2s ^{1}S - 1snp ^{1}P spectral series are fitted with polynomial function of 4-th order; the fit-curve is added in the figure. The data for 1s3p ^{1}P - 1snd ^{1}D spectral series are fitted with polynomial function of 3-th order. The polynomial function permits the interpolation and extrapolation to obtain electron impact width for other lines within the series where there is a lack of the data to perform the calculations. The analytical function up to 4-th order is given below:

$$W(n) = a_4 n^4 + a_3 n^3 + a_2 n^2 + a_1 n + a_0 \quad (7)$$

where W(n) is the full width at half-maximum in angular frequency units. The corresponding coefficients for the three series are given in Table 2.

As an example of astrophysical significance of obtained results Stark and Doppler line widths for B IV 2s ^{1}S - 2p ^{1}P^o (λ = 4492.1 Å) spectral line are compared in Fig. 2, as a function of Rosseland optical depth (τ) for an atmospheric model (Wesemael 1981) of helium rich DO white dwarf with



Figure 1. Principal quantum number dependence of the electron-impact width (in angular frequency units) for multiplets within 1s^2 ^{1}S - 1snp ^{1}P^o, 1s2s ^{1}S - 1snp ^{1}P^o and 1s3p ^{1}P^o - 1snd ^{1}D spectral series at an electron density of 1.10^{17} cm^{-3} and T = 200 000 K.



Figure 2. Stark and Doppler widths for B IV 2s ^{1}S - 2p ^{1}P^o (λ = 4492.1 Å) spectral line as a function of Rosseland optical depth (τ). Stark and Doppler widths are shown for an atmospheric model (Wesemael 1981) with effective temperatures T$_{eff}$ = 80 0000 K, and log g = 8.

the effective temperatures T$_{eff}$ = 80 0000 K, and log g = 8. One can see that Stark broadening dominates in the large part of the considered atmosphere.

4 CONCLUSIONS

New Stark broadening parameters for 157 multiplets of B IV have been calculated within the frame of the semi-classical perturbation formalism. The results for Stark broadening parameters of boron lines could be applicable for the adequate interpretation of the observed spectra in astrophysics, astrochemistry, and cosmology, in technological plasma research, for thermonuclear reaction devices, and for laser produced plasma investigation. We also note that the Stark broadening data obtained in the present research, will be in-

Table 1. This table gives electron-, proton-, and helium ion-impact broadening parameters for B IV lines. Calculated wavelength of the transitions (in Å) and parameter C are also given. This parameter, when divided with the corresponding Stark width, gives an estimate for the maximal pertuber density for which the line may be treated as isolated. The Stark broadening parameters for 157 B IV lines are available in its entirety in machine-readable form in the online journal as additional data. Results for perturber density of 10^{14} cm^{-3} are in Table 1, and for 10^{17} cm^{-3} - 10^{22} cm^{-3} in Tables 2-7. In all Tables temperatures are from 20 000 to 500 000 K. A positive shift is towards the red part of the spectrum. Data for 36 multiplet for electron density of 10^{17} cm^{-3} are from Dimitrijević et al. (2014) and here are added in order to complete Stark broadening data for B IV. A portion, for a perturber density of 10^{17} cm^{-3} is shown here for guidance regarding its form and content.

PERTURBER DENSITY = 1E+17cm-3							
TRANSITION	T(K)	ELECTRONS WIDTH(A)	SHIFT(A)	PROTONS WIDTH(A)	SHIFT(A)	IONIZED WIDTH(A)	HELIUM SHIFT(A)
SINGLETS							
B IV 2s-2p	20000.	0.171	-0.930E-02	0.354E-03	-0.208E-02	0.565E-03	-0.204E-02
4492.1 A	50000.	0.110	-0.646E-02	0.206E-02	-0.468E-02	0.252E-02	-0.437E-02
C= 0.45E+21	100000.	0.818E-01	-0.696E-02	0.466E-02	-0.692E-02	0.476E-02	-0.617E-02
	200000.	0.623E-01	-0.687E-02	0.835E-02	-0.924E-02	0.744E-02	-0.774E-02
	300000.	0.538E-01	-0.673E-02	0.101E-01	-0.103E-01	0.878E-02	-0.864E-02
	500000.	0.454E-01	-0.644E-02	0.122E-01	-0.117E-01	0.105E-01	-0.979E-02
B IV 2s-3p	20000.	0.336E-02	-0.199E-03	0.137E-03	-0.230E-03	0.151E-03	-0.201E-03
380.9 A	50000.	0.233E-02	-0.189E-03	0.319E-03	-0.375E-03	0.303E-03	-0.316E-03
C= 0.20E+18	100000.	0.181E-02	-0.181E-03	0.471E-03	-0.457E-03	0.402E-03	-0.386E-03
	200000.	0.143E-02	-0.163E-03	0.626E-03	-0.546E-03	0.513E-03	-0.462E-03
	300000.	0.126E-02	-0.145E-03	0.765E-03	-0.598E-03	0.573E-03	-0.499E-03
	500000.	0.107E-02	-0.125E-03	0.909E-03	-0.655E-03	0.675E-03	-0.552E-03

Table 2. Coefficients of the fitting polynomial functions for three spectral series. The first column presents the corresponding transition, the next 5 columns give the values of the coefficients in the fitting polynomial functions.

Transition	a_0	a_1	a_2	a_3	a_4
$1s^2\ ^1S$ - $1snp\ ^1P^o$	-4.2421E+11	5.2977E+11	-2.33982E+11	3-8929E+10	7.37826E+8
$1s2s\ ^1S$ - $1snp\ ^1P^o$	1.06075E+11	-1.54335E+9	-3.69292E+10	7.9373E+9	2.48152E+9
$1s3p\ ^1P^o$ - $1snd\ ^1D$	1.00003E+12	-1.91806E+11	-1.26934E+11	4.03773E+10	

serted in the STARK-B database (Sahal-Bréchot et al. 2012; Sahal-Bréchot et al. 2015a, 2010, 2015b), which is a part of Virtual Atomic and Molecular Data Center (VAMDC - (Dubernet et al. 2010; Rixon et al. 2011). Additionally, it has been confirmed that the behaviour of Stark broadening parameters within B IV spectral series is regular, enabling the interpolation and extrapolation of new data.

Also, an example of the significance of Stark broadening mechanism for BIV lines in DO white dwarf atmospheres is given.

ACKNOWLEDGMENTS

This work is a part of the project 176002 "Influence of collisional processes on astrophysical plasma line shapes" supported by the Ministry of Education, Science and Technological Development of Serbia. It is also the result of Short term Scientific Mission CM0805 within COST programme on The Chemical Cosmos: Understanding Chemistry in Astronomical Environments. Partial financial support from project No 6572-20 Technical University - Sofia within Euroatom Programme is also acknowledged. This work has been supported as well, by the Paris Observatory, the CNRS and the PNPS (Programme National de Physique Stellaire, INSU-CNRS).

REFERENCES

Barnard A. J., Cooper J., Smith E. W., 1974, J. Quant. Spectrosc. Radiat. Transfer, 14, 1025

Bates D. R., Damgaard A., 1949, Philos. Trans. R. Soc. London A, 242, 101

Blagojević, B., Popović, M. V., Konjević, N., Dimitrijević, M. S., 1999, J. Quant. Spectrosc. Radiat. Transfer, 61, 361

Boiko, V. A, Faenov, A. Ya., Hahalin, S. Ya., Pikuz, S. A., Shilov, K. A., Skobelev, I. Yu., 1983, J. Phys. B, 16, L77

Cunha, K., Smith, V. V., Lambert, D. L., 1999, ApJ, 519, 844

Cunha, K., Lambert, D. L., Lemke, M., Gies, D. R., Roberts, L. C., 1997, ApJ, 478, 211

Delbourgo-Salvador, P., Vangioni-Flam, E., 1993, in:

Prantzos et al. (Eds.), Origin and Evolution of Elements, Cambridge University Press, Cambridge, p. 52
Dimitrijević M. S., Sahal-Bréchot S., 1984, J. Quant. Spectrosc. Radiat. Transfer, 31, 301
Dimitrijević M. S., Sahal-Bréchot S., 1994, Phys. Scr., 49, 661
Dimitrijević M. S., Sahal-Bréchot S., 1996, Phys. Scr., 54, 50
Dimitrijević M. S., Sahal-Bréchot S., Bommier V., 1991, A&AS, 89, 581
Dimitrijević M. S., Jovanović P., Simić, Z., 2003a, A&A, 410, 735
Dimitrijević M. S., Ryabchikova T., Popović L. Č., Shulyak D., Tsymbal, V., 2003b, A&A, 404, 1099
Dimitrijević M. S., Dačić M., Cvetković Z., Simić Z., 2004, A&A, 425, 1147
Dimitrijević M. S., Ryabchikova T., Popović L. Č., Shulyak D., Khan S., 2005, A&A, 435, 1191
Dimitrijević M. S., Kovačević A., Simić Z., Sahal-Bréchot S., 2011, Baltic Astronomy, 20, 580
Dimitrijevic, M. S., Christova, M., Simić, Z., Kovačević, A., Sahal-Bréchot, S., 2014, Adv. Space Res., 54, 1195
Dubernet M. L., Boudon V., Culhane J. L., Dimitrijević M. S., Fazliev A. Z. et al., 2010, J. Quant. Spectrosc. Radiat. Transfer, 111, 2151, http://www.vamdc.org
Dufour P., Ben Nessib N., Sahal-Bréchot S., Dimitrijević M. S., 2011, Baltic Astronomy, 20, 511
Duncan D. K., Peterson R. C., Thorburn J. A., Pinsonneault M. H., 1998, **ApJ**, 499, 871
Duncan, D. K., Primas, F., Rebull, L. M., Boesgaard, A. M., Deliyannis, Constantine P., Hobbs, L. M., King, J. R., Ryan, S. G., 1997, ApJ, 488, 338
Fleurier C., Sahal-Bréchot S., Chapelle J., 1977, J. Quant. Spectrosc. Radiat. Transfer, 17, 595
Griem H. R., 1974, Spectral line Broadening by Plasmas. McGraw-Hill, New York
Hamdi R., Ben Nessib N., Milovanović N., Popović L. Č., Dimitrijević M. S., Sahal-Bréchot S., 2008, MNRAS, 387, 871
Iglesias, E., Griem, H., Welch, B., Weaver, J., 1998, Astrophys. Space Sci., 256, 327
Källne, E., Källne, J., Pradhan, A. K., 1983, Phys, Rev. A, 22, 1476
Kernahan, J. A., Pinnington, E. H., Livingston, A. E., Irwin, D. J. G., 1975, Phys. Scr., 12, 319
Kolk, K.-H., König, R., Kunze, H. -J., 1986, Phys. Rev. A, 33, 747
Kramida, A. E., Ryabtsev, A. N., Ekberg, J. O., Kink, I., Mannervik, S., Martinson, I., 2008, Phys. Scr., 78, 025302
Lanz T., Dimitrijević M. S., Artru M. C., 1988, A&A, 192, 249
Larbi-Terzi N., Sahal-Bréchot S., Ben Nessib N., Dimitrijević M. S., 2012, MNRAS, 423, 766
McKenzie, D. L. & Landecker, P. B., 1982, ApJ, 259, 372
Milovanović N., Dimitrijević M. S., Popović L. Č., Simić, Z., 2004, A&A, 417, 375
Nicolosi, P., Garifo, L., Jannitti, E., Malvezzi, A. M., Tondello, G., 1978, Nuovo Cimento B, 48, 133
Oertel G. K., Shomo L. P., 1968, ApJS, 16, 175
Popović L. Č., Dimitrijević M. S., Ryabchikova T., 1999a, A&A, 350, 719
Popović L. Č., Dimitrijević M. S., Tankosić D., 1999b, A&AS, 139, 617
Popović L. Č., Milovanović N., Dimitrijević M. S., 2001a, A&A, 365, 656
Popović L. Č., Simić S., Milovanović N., Dimitrijević M. S., 2001b, ApJS, 135, 109
Proffitt, C. R., Quigley, M. F., 2001, ApJ, 548, 429
Proffitt, C. R., Jönsson, P., Litzén, U., Pickering, J. C., Wahlgren, G. M., 1999, ApJ, 516, 342
Rixon G., Dubernet M. L., Piskunov, N., Walton, N., Mason, N., Le Sidaner, P., Schlemmer, S., Tennyson, J., et al., 2011 7^{th} International Conference on Atomic and Molecular Data and their Applications -ICAMDATA-2010, AIP Conf. Proc. 1344, 107
Sahal-Bréchot S., 1969a, A&A, 1, 91
Sahal-Bréchot S., 1969b, A&A, 2, 322
Sahal-Bréchot S., 1974, A&A, 35, 319
Sahal-Bréchot S., 1991, A&A, 245, 322
Sahal-Brechot, S., J. Phys.: Conf. Ser., 257, 012028
Sahal-Bréchot S., Dimitrijević M. S., Ben Nessib N., 2014, Atoms, 2, 225
Sahal-Bréchot, S., Dimitrijević, M.S., Moreau N., 2012, J. Phys.: Conf. Ser., 397, 012019
Sahal-Bréchot S., Dimitrijević M. S., Moreau N. 2015a, STARK-B database, [online]. Available: http://stark-b.obspm.fr [June 19, 2015]. Observatory of Paris, LERMA and Astronomical Observatory of Belgrade
Sahal-Bréchot S., Dimitrijević M. S., Moreau N., Ben Nessib N., 2015b, Phys. Scripta, 50, 054008
Schramm, D. N., 1993, in: Prantzos et al. (Eds.), Origin and Evolution of the Elements, Cambridge University Press, Cambridge, p. 112
Shore B. W., Menzel D., 1965, ApJS, 12, 187
Simić Z., Dimitrijević M. S., Popović L. Č., Dačić M., 2005a, J. Appl. Spectrosc., 72, 443
Simić Z., Dimitrijević M. S., Milovanović N., Sahal-Bréchot S., 2005b, A&A, 441, 391
Simić Z., Dimitrijević M. S., Popović L. Č., Dačić M., 2006, New Astron., 12, 187
Simić Z., Dimitrijević M. S., Kovačević A., 2009, New Astron. Rev., 53, 246
Simić Z., Dimitrijević M. S., Sahal-Bréchot S., 2013, MNRAS, 432, 2247
Simić Z., Dimitrijević M. S., Popović L. Č., 2014, Adv. Space Res., 54, 1231
Tankosić D., Popović L. Č., Dimitrijević M. S., 2003, A&A, 399, 795
TFR Group, Doyle, J. G., Schwob, J. L., 1982, J. Phys. B, 15, 813
Thomas, D., Schramm, D. N., Olive, K. A., Fields, B. D., 1993, ApJ, 406, 569
Van Regemorter H., Hoang Binh Dy, Prud'homme M., 1979, J. Phys. B, 12, 1073
Vangioni-Flam E., Cassé M., 1999, Astrophys. Space Sci., 265, 77
Vangioni-Flam E., Cass M., Audouze J., 2000, Phys. Rep. 333-334, 365
Venn K. A., Brooks A. M., Lambert, D. L., Lemke, M., Langer, N., Lennon, D. J., Keenan, F. P., 2002, ApJ, 565, 571
Wang, J. S., Griem, H. R., Huang, Y. W., Böttcher, F., 1992, Phys. Rev. A, 45, 4010

Wesemael F., 1981, ApJS, 4**5, 177**

This paper has been typeset from a TEX/ LATEX file prepared by the author.